



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Regis...]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Bool

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 0

1: J Pharm Pharmacol. 1988 Sep;40(9):627-33.

Links

Properties of a semicarbazide-sensitive amine oxidase in human umbilical artery.

Precious E, Lyles GA.

Department of Pharmacology and Clinical Pharmacology, University of Dundee, Ninewells Hospital, UK.

The metabolism of some aromatic amines by amine oxidase activities in human umbilical artery homogenates has been studied. The inhibitory effects of clorgyline showed that 5-hydroxytryptamine (5-HT) and tryptamine, 1 mM, were predominantly substrates for monoamine oxidase (MAO) type A, whereas MAO-A and B were both involved in the metabolism of beta-phenylethylamine (PEA), 100 microM, and tyramine, 1 mM. About 20-30% of tyramine and PEA metabolism was resistant to 1 mM clorgyline, but sensitive to inhibition by semicarbazide, 1 mM, indicating the presence of a semicarbazide-sensitive amine oxidase (SSAO). Benzylamine, 1 mM, appeared to be metabolized exclusively by SSAO with a K_m (161 microM) at pH 7.8 similar to that found for SSAO in other human tissues. Tyramine and PEA were relatively poor substrates for SSAO, with very high apparent K_m values of 17.6 and 13.3 mM, respectively, when determined in the presence of clorgyline, 10^{-3} M, added to inhibit any metabolism of those amines by MAO activities. However, kinetic studies with benzylamine indicated that clorgyline, 10^{-3} M, also appears to inhibit SSAO competitively such that the true K_m values for tyramine and PEA may be about 60% of those apparent values given above. No evidence for the metabolism of 5-HT or tryptamine by SSAO was obtained. The aliphatic amine methylamine was recently shown to be a specific substrate for SSAO in umbilical artery homogenates. We have used benzylamine and methylamine as SSAO substrates in histochemical studies to localize SSAO in tissue sections. (ABSTRACT TRUNCATED AT 250 WORDS)

PMID: 2907029 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort By Send to

Related Links

- Further studies on the metabolism of methylamine by semicarbazide-sensitive amine oxidase activities in human plasma, umbilical artery and rat aorta. [J Pharm Pharmacol. 1990]
- Deamination of methylamine by semicarbazide-sensitive amine oxidase in human umbilical artery and rat aorta. [Biochem Pharmacol. 1988]
- Amine oxidase activities in brown adipose tissue of the rat: identification of semicarbazide-sensitive (clorgyline-resistant) activity at the fat cell membrane. [J Pharm Pharmacol. 1984]
- Semicarbazide-sensitive amine oxidase (SSAO) of the rat aorta. Interactions with some naturally occurring amines and their structure. [Biochem Pharmacol. 1989]
- The metabolism of aminoacetone to methylglyoxal by semicarbazide-sensitive amine oxidase in human umbilical artery. [Biochem Pharmacol. 1992]

See all Related Articles...

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)
Department of Health & Human Services
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Regis]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 0 X

1: J Auton Pharmacol. 1991 Oct;11(5):323-35.

Links

Effect of benzylamine and its metabolites on the responses of the isolated perfused mesenteric arterial bed of the rat.

Elliott J, Callingham BA.

Department of Pharmacology, University of Cambridge, UK.

1. Semicarbazide-sensitive amine oxidase (SSAO) is an enzyme activity which can be found in the plasma membrane of rat vascular smooth muscle cells. We have investigated the possibility that the products of deamination by this enzyme, namely ammonia, hydrogen peroxide and the aldehyde, may be important in the modulation of the responses of vascular smooth muscle to extracellular stimuli. 2. The isolated perfused mesenteric arterial bed of the rat was used and dose-pressure response curves (DRC) to bolus injections of adrenaline (Ad) or ATP were plotted by non-linear curve fitting. The relaxant effects of carbachol (CCh), which releases endothelium dependent relaxing factor (EDRF), were studied by co-administering CCh with Ad. The effects of including the preferred SSAO substrate, benzylamine (BZ; 25 microM), in the perfusion fluid throughout the experiment and of inhibition of SSAO by treatment of rats with (E)-2-(3',4'-dimethoxyphenyl)-3-fluoroallylamine (MDL 72145; 1 mg kg⁻¹) 1 h before dissection, have been studied. 3. Neither BZ nor SSAO inhibition affected the DRC to ATP. BZ shifted Ad responses to the left, inhibition of SSAO increased this shift indicating that the amine, but not its metabolites, were responsible for the potentiation of the responses to Ad. DRC to CCh showed a shift to the left and a significant decrease in the Hill slope with BZ, indicative of a potentiation of low doses of CCh more than high doses. Inhibition of SSAO prevented this change and so the metabolites of BZ deamination appeared to be involved in the potentiation. 4. Ammonia generated by SSAO may contribute to the production of EDRF or hydrogen peroxide may sensitize guanylate cyclase to stimulation by EDRF and so explain these findings.

PMID: 1748692 [PubMed - indexed for MEDLINE]

Related Links

- ▷ The influence of amine metabolizing enzymes on the pharmacology of tyramine in the isolated perfused mesenteric arterial bed of the rat. [Br J Pharmacol. 1989]
- ▷ Acrolein generation stimulates hypercontraction in isolated human blood vessels. [Toxicol Appl Pharmacol. 2006]
- ▷ Metabolism of amines in the isolated perfused mesenteric arterial bed of the rat. [Br J Pharmacol. 1989]
- ▷ Semicarbazide-sensitive amine oxidase (SSAO) of the rat aorta. Interactions with some naturally occurring amines and their structure. [Biochem Pharmacol. 1989]
- ▷ The role of semicarbazide-sensitive amine oxidase with a high affinity for benzylamine (Bz. SSAO) in the catabolism of histamine in the mesenteric arterial bed of the rat. [Agents Actions. 1994]

See all Related Articles...

Display AbstractPlus Show 20 Sort By Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)


[Home](#) [Browse](#) [Search](#) [My Settings](#) [Alerts](#) [Help](#)


Quick Search Title, abstract, keywords Author e.g.
 search tips Journal/book title Volume Issue Page

Biochemical Pharmacology
 Volume 38, Issue 9, 1 May 1989, Pages 1507-1515

Abstract

 Add to my Quick Links  Cited By  E-mail Article  Save as Citation Alert  Export Citation  Citation Feed

doi:10.1016/0006-2952(89)90191-3  Cite or Link Using DOI
 Copyright © 1989 Published by Elsevier Science Inc.

 Request Permission

[View Record in Scopus](#)
[Cited By in Scopus \(18\)](#)

Semicarbazide-Sensitive Amine Oxidase (SSAO) of the rat aorta

Interactions with some naturally occurring amines and their structural analogues

Jonathan Elliott, Brian A. Callingham and Dennis F. Sharman

Department of Pharmacology, University of Cambridge, Hills Road, Cambridge, CB2 2QD, U.K.

Received 30 August 1988; accepted 21 November 1988. Available online 15 November 2002.

Abstract

The influence of a number of naturally occurring amines and their structural analogues has been examined on the metabolism of radiolabelled benzylamine (BZ) by the membrane bound semicarbazide-sensitive amine oxidase (SSAO) of the rat aorta. Only primary monoamines were effective in reducing the deamination of BZ. In the phenylethylamine series, addition of hydroxyl groups to the benzene ring decreased their potency as inhibitors while addition of a hydroxyl group at the β position increased the inhibitory potency. Stereoselectivity of action was shown with octopamine, the L-isomer being the more active form. Kinetic analysis of these interactions showed predominantly competitive inhibition and kynuramine had the lowest K_i of 5.4 μ M. The aliphatic monoamines, isoamylamine and isobutylamine both competed with BZ. 5-Hydroxytryptamine (5-HT) was the only amine that inhibited non-competitively. Direct evidence for metabolism by SSAO of some of the competing amines such as isoamylamine, phenylethylamine, tyramine and tryptamine was obtained by fluorimetric or radio-chemical assays. The inhibitors clorgyline and (E)-2-(3',4'-dimethoxyphenyl)-3-fluoroallylamine (MDL 72145) were used to characterise the amine oxidase activity responsible for the deamination. Octopamine and phenylethanolamine (PeOH) were not SSAO substrates and inhibited BZ metabolism in the fluorimetric assay. It is possible that the activity of SSAO is controlled by octopamine released

from sympathetic nerve endings or 5-HT released from platelets.

Biochemical Pharmacology

Volume 38, Issue 9, 1 May 1989, Pages 1507-1515

[Home](#) [Browse](#) [Search](#) [My Settings](#) [Alerts](#) [Help](#)



[About ScienceDirect](#) | [Contact Us](#) | [Terms & Conditions](#) | [Privacy Policy](#)

Copyright © 2007 Elsevier B.V. All rights reserved. ScienceDirect® is a registered trademark of Elsevier B.V.



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Regis

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 0

1: Biochem Pharmacol. 1989 May 1;38(9):1507-15.

ELSEVIER Links
FULL-TEXT ARTICLE

Semicarbazide-sensitive amine oxidase (SSAO) of the rat aorta. Interactions with some naturally occurring amines and their structural analogues.

Elliott J, Callingham BA, Sharman DF.

Department of Pharmacology, University of Cambridge, U.K.

The influence of a number of naturally occurring amines and their structural analogues has been examined on the metabolism of radiolabelled benzylamine (BZ) by the membrane bound semicarbazide-sensitive amine oxidase (SSAO) of the rat aorta. Only primary monoamines were effective in reducing the deamination of BZ. In the phenylethylamine series, addition of hydroxyl groups to the benzene ring decreased their potency as inhibitors while addition of a hydroxyl group at the beta position increased the inhibitory potency. Stereoselectivity of action was shown with octopamine, the L-isomer being the more active form. Kinetic analysis of these interactions showed predominantly competitive inhibition and kynuramine had the lowest K_i of 5.4 μM . The aliphatic monoamines, isoamylamine and isobutylamine both competed with BZ. 5-Hydroxytryptamine (5-HT) was the only amine that inhibited non-competitively. Direct evidence for metabolism by SSAO of some of the competing amines such as isoamylamine, phenylethylamine, tyramine and tryptamine was obtained by fluorimetric or radiochemical assays. The inhibitors clorgyline and (E)-2-(3',4'-dimethoxyphenyl)-3-fluoroallylamine (MDL 72145) were used to characterise the amine oxidase activity responsible for the deamination. Octopamine and phenylethanolamine (PeOH) were not SSAO substrates and inhibited BZ metabolism in the fluorimetric assay. It is possible that the activity of SSAO is controlled by octopamine released from sympathetic nerve endings or 5-HT released from platelets.

PMID: 2719723 [PubMed - indexed for MEDLINE]

Related Links

- ▷ Deamination of methylamine by semicarbazide-sensitive amine oxidase in human umbilical artery and rat aorta. [Biochem Pharmacol. 1988]
- ▷ Properties of a semicarbazide-sensitive amine oxidase in human umbilical artery. [J Pharm Pharmacol. 1988]
- ▷ Inhibition of rat aorta semicarbazide-sensitive amine oxidase by 2-phenyl-3-haloallylamines and related compounds. [Biochem Pharmacol. 1987]
- ▷ Several aspects on the amine oxidation by semicarbazide-sensitive amine oxidase (SSAO) in rat aorta. [Neurotransm Suppl 1994]
- ▷ An allylamine derivative (MDL 72145) with potent irreversible inhibitory actions on rat aorta semicarbazide-sensitive amine oxidase. [Biochem Pharmacol. 1985]

See all Related Articles...

Display AbstractPlus Show 20 Sort By Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Regis

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Bool

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 0

1: J Pharm Pharmacol. 1989 Feb;41(2):97-100.

Links

The enhanced daily excretion of urinary methylamine in rats treated with semicarbazide or hydralazine may be related to the inhibition of semicarbazide-sensitive amine oxidase activities.

Lyles GA, McDougall SA.

Department of Pharmacology and Clinical Pharmacology, University of Dundee, Ninewells Hospital and Medical School, Scotland, UK.

The effects of amine oxidase inhibitors upon the daily urinary excretion of monomethylamine (MMA), dimethylamine (DMA), trimethylamine (TMA) and ammonia in the rat have been examined. Administration of hydralazine (5 mg kg⁻¹) or semicarbazide (100 mg kg⁻¹), drugs which irreversibly inhibit semicarbazide-sensitive amine oxidases (SSAO) but not monoamine oxidase (MAO), enhanced MMA excretion by around three- to six-fold above pretreatment levels, whereas no effect of pargyline (25 mg kg⁻¹), a selective irreversible inhibitor of MAO was found. No apparent changes in DMA or TMA excretion in response to drug-treatment were observed. Ammonia excretion also was generally unchanged except for an apparent marked increase (approximately four-fold) over the 24 h following semicarbazide, a result which might be explained if ammonia is a degradation product of semicarbazide metabolism in the rat. With recent evidence that MMA is a substrate in-vitro for SSAO activities, results here may indicate that SSAO or related enzymes are involved in endogenous MMA turnover.

PMID: 2568436 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort By Send to

Related Links

- ▶ Deamination of methylamine by semicarbazide-sensitive amine oxidase in human umbilical artery and rat aorta. [Biochem Pharmacol. 1988]
- ▶ The interaction of hydralazine with a semicarbazide-sensitive amine oxidase in brown adipose tissue of the rat. Its use as a radioactive ligand for the enzyme. [Biochem J. 1985]
- ▶ Hydralazine is an irreversible inhibitor of the semicarbazide-sensitive, clorgyline-resistant amine oxidase in rat aorta homogenates. [J Pharm Pharmacol. 1982]
- ▶ Glucose handling in streptozotocin-induced diabetic rats is improved by tyramine but not by the amine oxidase inhibitor semicarbazide. [Life Sci. 2005]
- ▶ Methylamine metabolism to formaldehyde by vascular semicarbazide-sensitive amine oxidase. [Toxicology. 1992]

See all Related Articles...

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

Department of Health & Human Services

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Regis]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for Go Clear

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 0

1: J Pharm Pharmacol. 1985 Sep;37(9):637-43.

Links

Vascular smooth muscle cells: a major source of the semicarbazide-sensitive amine oxidase of the rat aorta.

Lyles GA, Singh I.

Several methods have been used to study the distribution of the semicarbazide-sensitive amine oxidase (SSAO) within the wall of the rat aorta. After separation of the smooth muscle-containing layers of the tunica media from the connective tissue of the tunica adventitia, much higher specific enzyme activity (measured with 1 microM benzylamine) was found in homogenates of the media than of adventitia. Similar results were obtained for MAO-A (with 1 mM 5-HT as substrate). SSAO activity was also considerably higher in homogenates of cells (predominantly smooth muscle) isolated from medial tissue by enzymatic dissociation with collagenase and elastase compared with homogenates of cells (mostly of connective tissue origin) from the adventitia. Histochemical staining resulting from SSAO activity (with benzylamine as substrate) occurred predominantly and intensely over the tunica media in rat aortic sections, although some occasional staining of adventitial sites was also observed. Staining was prevented by the SSAO inhibitors hydroxylamine (1 microM) and semicarbazide (1 mM), but not by the MAO inhibitor, clorgyline (1 mM). These results indicate that SSAO is associated predominantly, although not exclusively, with the smooth muscle cells in the rat aorta. Our findings that beta-aminopropionitrile (BAPN) is a reversible, competitive inhibitor (K_i around 2×10^{-4} M) of SSAO, in contrast to the irreversible inhibition of the connective tissue lysyl oxidase by BAPN reported by others, provides further evidence that these enzymes are not identical.

PMID: 2867184 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort By Send to

Related Links

- ▶ Aminoacetone metabolism by semicarbazide-sensitive amine oxidase in rat aorta. [Biochem Pharmacol. 1995]
- ▶ Properties of a semicarbazide-sensitive amine oxidase in human umbilical artery. [J Pharm Pharmacol. 1988]
- ▶ Semicarbazide-sensitive amine oxidase activity in rat aortic cultured smooth muscle cells. [J Neural Transm Suppl. 1990]
- ▶ Amine oxidase activities in brown adipose tissue of the rat: identification of semicarbazide-sensitive (clorgyline-resistant) activity at the fat cell membrane. [J Pharm Pharmacol. 1984]
- ▶ Monoamine oxidase activities in dissociated cell fractions from rat skeletal muscle. [J Pharm Pharmacol. 1986]

See all Related Articles...

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Regis]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for Go Clear

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 1

1: Prog Brain Res. 1995;106:293-303.

Links

Substrate-specificity of mammalian tissue-bound semicarbazide-sensitive amine oxidase.

Lyles GA.

Department of Pharmacology and Clinical Pharmacology, University of Dundee, Ninewells Hospital and Medical School, UK.

Although the existence of a membrane-bound (probably plasmalemmal) semicarbazide-sensitive amine oxidase (SSAO) is well established in various mammalian tissues, and especially within vascular smooth muscle, its importance and the possible consequences of its metabolism of certain physiological and xenobiotic amines in vivo are under continuing investigation. In this respect, there are major species-related differences in substrate specificity determined in vitro, not only towards the synthetic amine benzylamine, but also towards some other aromatic amines (e.g. tyramine, tryptamine, 2-phenylethylamine, dopamine, histamine) which are possible endogenous substrates. Inhibition of SSAO can potentiate the pharmacological activity of some amines in isolated tissue (e.g. blood vessel) preparations from some species. Recent evidence has accumulated that SSAO may also be involved in metabolizing endogenous aliphatic amines such as methylamine and aminoacetone, focussing attention on the fact that the aldehyde products (formaldehyde and methylglyoxal, respectively) are potentially cytotoxic agents. Indeed, SSAO has been implicated in experimental models of cardiovascular toxicity involving conversion of the industrial aliphatic amine allylamine to acrolein. In summary, metabolism by SSAO may reduce the physiological/pharmacological effects of some amines, but the resulting metabolites (aldehydes, H₂O₂) may also have important actions.

PMID: 8584666 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort By Send to

Related Links

- ▶ Properties of mammalian tissue-bound semicarbazide-sensitive amine oxidase: possible clues to its physiological function [Neurotransm Suppl. 1994]
- ▶ Mammalian plasma and tissue-bound semicarbazide-sensitive amine oxidases: biochemical, pharmacological and toxicological aspects. [Int J Biochem Cell Biol. 1996]
- ▶ Several aspects on the amine oxidation by semicarbazide-sensitive amine oxidase (SSAO) in the brain [Neurotransm Suppl. 1994]
- ▶ Some aspects of the pathophysiology of semicarbazide-sensitive amine oxidase enzymes. [Prog Brain Res. 1995]
- ▶ The metabolism of aminoacetone to methylglyoxal by semicarbazide-sensitive amine oxidase in human umbilical artery [Biochem Pharmacol. 1992]

See all Related Articles...

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Register]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 0

1: Biochem Pharmacol. 1983 Sep 1;32(17):2515-21.

ELSEVIER Links
FULL-TEXT ARTICLE

Inhibitory actions of hydralazine upon monoamine oxidizing enzymes in the rat.

Lyles GA, Garcia-Rodriguez J, Callingham BA.

The inhibition by hydralazine of the clorgyline-resistant amine oxidase (CRAO) and monoamine oxidase (MAO) activities in various rat tissues has been studied. Hydralazine was a potent, time-dependent inhibitor of rat heart CRAO activity in vitro. The inhibition was not reversed by dialysis for 18 hr at 4 degrees, and only partially reversed by dialysis at 37 degrees. Dialysis at 4 degrees in the presence of pyridoxal phosphate (10(-4) M) also did not reverse the inhibition. Ex vivo inhibition of CRAO was found in heart and aorta homogenates in a dose-dependent manner after administration of hydralazine (1-40 mg/kg i.p.) to rats. In contrast, MAO-A activity was unaffected or, in some cases, significantly increased in these tissue homogenates from drug-treated animals. However, in vitro inhibition by hydralazine of both MAO-A and B activities of rat liver mitochondrial fractions was found, and these effects were fully reversible by dialysis for 18 hr at 4 degrees. Inhibition of MAO-A was competitive (K_i of 2.5×10^{-6} M), while inhibition of MAO-B showed complex mixed non-competitive kinetics. These results indicate that hydralazine possesses different inhibitory properties towards the various amine oxidases in rat tissues, and these actions are discussed in relation to the clinical use of the drug as an anti-hypertensive agent.

PMID: 6615547 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort By Send to

Related Links

- ▶ The effect of DSP-4 (N-[2-chloroethyl]-N-ethyl-2-bromobenzylamine) on monoamine oxidase activities in tissues of the rat. [Pharm Pharmacol. 1981]
- ▶ In vitro and in vivo inhibition by benserazide of clorgyline-resistant amine oxidases in rat cardiovascular tissues. [Biochem Pharmacol. 1982]
- ▶ Monoamine oxidase-inhibiting properties of SR 95191, a new pyridazine derivative, in the rat: evidence for selective and reversible inhibition of monoamine oxidase type A in vivo but not in vitro. [Neurochem. 1988]
- ▶ Inhibition of monoamine oxidase A and B activities by imidazol(ine)/guanidine drugs, nature of the interaction and distinction from 12-imidazoline receptors in rat liver. [Br J Pharmacol. 1997]
- ▶ Neurochemical profile of moclobemide, a short-acting and reversible inhibitor of monoamine oxidase. [Pharmacol Ther. 1989]

See all Related Articles...

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Regis]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Bool

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 0

1: Biochem Pharmacol. 1983 Sep 1;32(17):2515-21.

ELSEVIER Links
FULL-TEXT ARTICLE

Inhibitory actions of hydralazine upon monoamine oxidizing enzymes in the rat.

Lyles GA, Garcia-Rodriguez J, Callingham BA.

The inhibition by hydralazine of the clorgyline-resistant amine oxidase (CRAO) and monoamine oxidase (MAO) activities in various rat tissues has been studied. Hydralazine was a potent, time-dependent inhibitor of rat heart CRAO activity in vitro. The inhibition was not reversed by dialysis for 18 hr at 4 degrees, and only partially reversed by dialysis at 37 degrees. Dialysis at 4 degrees in the presence of pyridoxal phosphate (10^{-4} M) also did not reverse the inhibition. Ex vivo inhibition of CRAO was found in heart and aorta homogenates in a dose-dependent manner after administration of hydralazine (1-40 mg/kg i.p.) to rats. In contrast, MAO-A activity was unaffected or, in some cases, significantly increased in these tissue homogenates from drug-treated animals. However, in vitro inhibition by hydralazine of both MAO-A and B activities of rat liver mitochondrial fractions was found, and these effects were fully reversible by dialysis for 18 hr at 4 degrees. Inhibition of MAO-A was competitive (K_i of 2.5×10^{-6} M), while inhibition of MAO-B showed complex mixed non-competitive kinetics. These results indicate that hydralazine possesses different inhibitory properties towards the various amine oxidases in rat tissues, and these actions are discussed in relation to the clinical use of the drug as an anti-hypertensive agent.

PMID: 6615547 [PubMed - indexed for MEDLINE]

Related Links

- ▶ The effect of DSP-4 (N-[2-chloroethyl]-N-ethyl-2-bromobenzylamine) on monoamine oxidase activities in tissues of the rat. [Pharmacol. 1981]
- ▶ In vitro and in vivo inhibition by benserazide of clorgyline-resistant amine oxidases in rat cardiovascular tissues. [Biochem Pharmacol. 1982]
- ▶ Monoamine oxidase-inhibiting properties of SR 95191, a new pyridazine derivative, in the rat: evidence for selective and reversible inhibition of monoamine oxidase type A in vivo but not in vitro. [Neurochem. 1988]
- ▶ Inhibition of monoamine oxidase A and B activities by imidazole/guanidine drugs, nature of the interaction and distinction from 12-imidazoline receptors in rat liver. [Pharmacol. 1997]
- ▶ Neurochemical profile of moclobemide, a short-acting and reversible inhibitor of monoamine oxidase. [Pharmacol. Ther. 1989]

See all Related Articles...

Display AbstractPlus Show 20 Sort By Send to

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)